Pur-Athlete®





An extruded high fat supplement derived from vegetable & flaxseed (source of Omega 3) and contains blood building nutrients (copper, cobalt, folic acid, Vitamin B12). In addition to improving skin, hair coat, hoof condition and muscle mass, research demonstrates that fat spares muscle glycogen. Pur-Athlete as a controlled NSC (sugars and starch) level of 17% max.

Analysis

Crude Protein (min.)	20.00 %	Phosphorus (actual)	0.60 %
Crude Fat (min.)	25.00 %	Sodium (actual)	0.25 %
Crude Fat (max.)	27.00 %	Vitamin A (min.)	13000 ^{I.U./kg}
Crude Fibre (max.)	4.00 %	Vitamin D3 (min.)	1300 ^{I.U./kg}
Calcium (actual)	0.70 %	Vitamin E (min.)	500 ^{I.U./kg}

Feeding rates

Administer 500 grams (per 500 kg body weight) orally with each meal, twice daily.

Characteristics

Unique, advanced feed formulation	Provides complete nutrient fortification required by performance horses.
25 - 27% fat content (vegetable oil)	 A more concentrated and efficient energy source than grains. Builds energy reserves and fuels superior performance. "Clean" burning fuel, which reduces metabolic waste and lactic acid build-up. Improves skin, hair, hoof condition and overall appearance.
Contains blood building nutrients (copper, cobalt, folic acid, Vitamin B12)	Builds cardiovascular capacity.
High protein	Builds muscle mass.
Extruded feed form	Higher digestibility.Ensures optimum feed intake.Mixes well with other sweet feeds, pellets and grains.
Contains Flaxseed (an excellent source of Omega-3)	Boosts immune function. Enhances anti-inflammatory properties and overall health.



Ingredients

Soybean Meal, DL-Methionine, L-Lysine, Corn Gluten Meal
Soya Oil, Flaxseed
Vitamin A, Vitamin D3, Vitamin E, Vitamin K, Riboflavin, Niacin, Thiamine, Vitamin B12, Biotin, Choline Chloride, Pantothenic Acid, Folic Acid, Vitamin B6
Salt (Sodium Chloride), Calcium Carbonate
Wheat shorts, Wheat germ

^{*} Ingredients are not in order of their inclusion rate.

- 1. Purina recommends analyzing your hay and balancing the ration accordingly.
- Feed clean and dust free hay in sufficient amount daily. Feeding poor quality hay or feeding inadequate amounts could cause colic and other digestive upsets.
- 3. Offer source of salt (NaCl) daily as well as plenty of fresh and clean water at all times.
- 4. Make any feed change gradually over a 7 to 10 days period.
- 5. Daily ration should be divided into 2 and pre-ferably 3 separate feedings or more.
- 6. Use a scale to weigh your feed so you know exactly how much your horse is being fed. Do not feed by volume.
- These recommendations are intended as a guide. They should be adjusted to variables of management, environment and individual needs. Consult your Purina Equine Consultant if necessary. (pam@cfs.ca)
- 8. Follow sound management practices along with optimal preventive veterinary medicine.

Caution

Directions for use must be carefully followed.

Feed is perishable. Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, abortion or death.