

# SUPPLEMENTS

#35820



# BMZ<sup>®</sup>

BMZ is a biotin, methionine and zinc (organic) supplement. A quality hoof care topdress formulated specifically to increase hoof growth and integrity. Methionine is interchangeable with cysteine, a major amino acid of keritin, which is the main protein in connective tissue, such as hoof wall. Research has demonstrated that hoof integrity is directly related to zinc content in the hoof wall.

## Analysis

<b>Biotin</b> (min.)	667 mg/kg
<b>Zinc</b> (actual)	5000 mg/kg

## Feeding rates

Administer ½ scoop (15 grams per 500 kg body weight) orally with each meal, twice daily. 30 gram scoop enclosed.

Note: Feeding vitamins other than vitamin A, D, E, riboflavin, pyridoxine and thiamin to horses may not have a beneficial effect.

## Characteristics

20 mg of biotin per dose	• 15 to 20 mg of biotin per day is scientifically proven to enhance hoof integrity and growth.
Contains D-L methionine	• Methionine is interchangeable with cysteine, a major amino acid of keritin, which is the main protein in connective tissue, such as hoof wall.
Contains zinc proteinate	• Chelation ensures enhanced digestibility and research has demonstrated that hoof integrity is directly related to hoof wall zinc content.
Contains dehydrated alfalfa meal as a carrier	• Enhanced palatability.
Manufactured in a plant with high standards of biosecurity and quality	• Consistent quality feed.
Manufactured in a 100% drug-free facility	• No risk of detrimental drug contamination. • Peace of mind.
Manufactured in Canada	• Minimizes transport cost. • Optimizes freshness.



## Ingredients

Fibre	Dehydrated Alfalfa Meal
Protein/Amino Acids	DL-Methionine
Starch	Ground Wheat
Fat	Soya Oil
Probiotic	
Vitamins	Biotin
Minerals	Organic Zinc
Multi-Attributes	Wheat shorts
Flavour/Binder	Flavouring

\* Ingredients are not in order of their inclusion rate.

1. Purina recommends analyzing your hay and balancing rations accordingly.
2. Feed clean and dust free hay in sufficient amounts daily. Feeding poor quality hay or feeding inadequate amounts could cause colic and other digestive upsets.
3. Offer source of salt (NaCl) daily as well as plenty of fresh and clean water at all times.
4. Make any feed change gradually over a 7 to 10 days period.
5. Daily ration should be divided into 2 and preferably 3 separate feedings or more.
6. Use a scale to weigh your feed so you know exactly how much your horse is being fed. Do not feed by volume.
7. These recommendations are intended as a guide. They should be adjusted to variables of management, environment and individual needs. Consult your CFS Purina Equine Specialist if necessary. 902-890-5354
8. Follow sound management practices along with optimal preventative veterinary medicine.

## Caution

Directions for use must be carefully followed.

Feed is perishable. Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, abortion or death.